



RRB

Section Controller

Railway Recruitment Board (RRB)

Volume - 1

Analytical and Mathematical Capability



INDEX

S No.	Chapter Title	Page No.
1	Number System	1
2	LCM & HCF	11
3	Surds and Indices	14
4	Simplification	17
5	Ratio, proportion and variation	21
6	Age	25
7	Average	26
8	Percentage	30
9	Profit and Loss	35
10	Discount and Dishonest shopkeeper	39
11	Mixture and Alligation	42
12	Partnership	47
13	Time Speed and Distance	49
14	Boat & Stream	54
15	Time and Work	56
16	Pipe and Cistern	60
17	Algebra	62
18	Linear Equations	68
19	Sequence and series	70
20	Lines and angle	74
21	Triangle	76
22	Circle	88
23	Quadrilaterals	96

INDEX

S No.	Chapter Title	Page No.
24	Polygon	102
25	Mensuration 2D	105
26	Mensuration 3D	117
27	Probability	126
28	Statistics	133
29	Data Sufficiency	142
30	Data Interpretation	146


1

CHAPTER

Number System



- **Number System:** A number system is a **method of representing and working with numbers** using a defined set of symbols and rules.




Types	Definition
Real Number	A real number is any number that can be represented on the number line. A real number is a number that includes all rational and irrational numbers and can be expressed as a point on the number line.
Rational Number	A rational number is a number that can be expressed in the form $\frac{p}{q}$, where p and q are integers and $q \neq 0$.

Irrational Number	An irrational number is a number that cannot be expressed in the form $\frac{p}{q}$, where p and q are integers and $q \neq 0$
Fraction	A fraction is a number that represents a part of a whole or a ratio of two quantities. It is written in the form $\frac{a}{b}$.
Integer	An integer is a whole number that can be positive, negative, or zero, and does not include any fractional or decimal part

Negative Integer	Negative integers are whole numbers with a negative sign, such as $-1, -2, -3, \dots$	Co-Prime Numbers	Two or more numbers are called co-prime (or relatively prime) if their only common factor (HCF) is 1. ➤ 1 is neither prime or composite.
Whole Number	A whole number is a non-negative integer, including zero.	Twin Prime Numbers	Twin numbers (twin primes) are a pair of prime numbers whose difference is exactly 2. ➤ 5 is only prime which is in 2 twin prime pairs. (3, 5) (5, 7) ➤ Sum of twin prime number (except 3 and 5) is always divisible by 12.
Natural number	Natural numbers are the numbers starting from 1 and increasing by 1 each time. 1, 2, 3, 4, ...	Decimal Number	A decimal number is a number that has a decimal point (.) and consists of a whole part and a fractional part. Ex- 3.5, 12.75 etc
Odd Number	An odd number is a natural number that is not divisible by 2 or in the form of $2n + 1$	Terminating Decimal	A terminating decimal is a decimal number that ends after a finite number of digits after the decimal point. Ex- 2.5, 0.75
Even Numbers	An even number is a natural number that is completely divisible by 2 or in the form of $2n$.	Non-Terminating Repeating Decimal	A non-terminating repeating decimal is a decimal number that does not end and in which one or more digits repeat continuously after the decimal point. Ex- 0. 333., 0. 1212
Prime number	A prime number is a natural number greater than 1 that has exactly two distinct factors: 1 and the number itself. 2, 3, 5, 7. ➤ 2 is the smallest prime number. ➤ 2 is only even prime number. ➤ All prime number (except 2 and 3) can be written in the form of $6n + 1$ or $6n + 5$ where n is natural number however, the converse is not true. ➤ (3, 5, 7) is only group of three primes, which are consecutive odds. ➤ 101 is smallest 3-digit prime number. ➤ 997 is largest 3-digit prime number.	Non-Terminating Non-Repeating Decimal	A non-terminating non-repeating decimal is a decimal number that does not end and does not repeat any digit or pattern after the decimal point. Ex- 1.1412, 3.14
Composite Number	A composite number is a natural number greater than 1 that has more than two factors. A composite number can be divided exactly by 1, itself, and at least one other number. 4, 6, 8, 9, 10, 12, 14... ➤ The smallest composite number is 4. ➤ The smallest odd composite number is 9 If a and b are any two odd primes then $a^2 + b^2$ and $a^2 - b^2$ is composite numbers.	Imaginary Number	An imaginary number is a number that can be written in the form $= bi$ b is a real number i is the imaginary unit, defined by

Complex Number	A complex number is a number that consists of two parts—a real part and an imaginary part—and can be expressed in the standard form $z = a + ib$ a is a real number, called the real part of z b is a real number, called the imaginary part of z i is the imaginary unit, defined by the property
----------------	---

Integer Number Line



How to check the given number is prime or not?

- To determine whether a number is prime, first find its square root and round it down to the nearest whole number. Then check whether the number is divisible by any prime number up to this value. If it is not divisible by any of them, the number is a prime number.

Between	Number or prime number
1-50	15
1-100	25
1-200	46

Ramanujan number

A Ramanujan number is a number that can be expressed as the sum of two positive cubes in two different ways. It is also known as the Hardy-Ramanujan number or taxi-cab number.

Smallest Ramanujan number = 1729

$$1729 = 1^3 + 12^3 = 9^3 + 10^3$$

Perfect number

A perfect number is a natural number that is equal to the sum of its proper divisors (that is, all its positive divisors excluding the number itself).

Ex: 4 is divisible by 1 and 2, so $1 + 2 = 3 \neq 4$; therefore, 4 is not a perfect number.

6 is divisible by 1, 2 and 3, so, $1 + 2 + 3 = 6 = 6$; therefore, 6 is a perfect number.

Key points

Even + Even = Even

Even $\times$ Even = Even

Even + Odd = odd

Even $\times$ Odd = Even

Odd + odd = Even

Odd $\times$ Odd = Odd

Sum Of 'n' odd number is odd, if n is odd.

Sum of 'n' odd number is even, if n is even.

Type 1: Question based on definitions



Ex: 173 is prime number or not?

Sol: The Square root of 173 is approximately 13.

The Prime number less than or equal to 13 are 2, 3, 5, 7, 11 and 13,

Since 173 is not divisible by any number, it is a prime number.

Ex: x, y and z are distinct prime number where $x < y < z$ if $x + y + z = 70$, then what is the value of z ?

Sol: Now, sum is 70, means at least one of the numbers is even. As we know, only one even prime number exists, and that is 2.

2 is also the smallest prime number.

Means $x = 2$

Now, $70 - 2 = 68 = y + z$

By substituting the values of different prime numbers, we obtain the result.

$y = 31$ and $x = 37$

Ex: How many composite numbers are there from 53 to 97?

Sol: If we find the total number of integers between 53 and 97 and then subtract the number of prime numbers between them, we obtain the number of composite numbers.

Total number = $97 - 53 + 1 = 45$ (+1 when both numbers are included)

Total prime number from 53 to 97 is 10

So, composite number = $45 - 10 = 35$

Ex: Which one of the following is TRUE?

- (A) All irrational numbers are real
- (B) All real numbers are irrational.
- (C) Rational numbers are not real.
- (D) Integers are not rational.

Sol: Irrational numbers form a subset of real numbers, so all irrational numbers are real.

Not all real numbers are irrational; rational numbers are also real.

Rational numbers are real numbers.

Integers are a subset of rational numbers since any integer can be written as $\frac{p}{q}$ (e.g., $5 = \frac{5}{1}$).

So, Answer is (A).

Special concept: based on sum of digits for special types of numbers

Number	Square	Sum of digits
11^2	121	3
111^2	12321	9
So, on		
111111111^2	12345678987654321	81

Ex: Each digit of a 9-digit number is 1. It is multiplied by itself. What is the sum of digit of the resulting number.

Sol- using the concept

$$111111111^2 = \gg 81$$

Type 2: Unit digit based

To find the units digit of an expression, consider only the unit digits of the numbers instead of evaluating the entire expression.

Unit digit of $(a + b)$ = unit digit of a + unit digit of b

Unit digit of $(a - b)$ = unit digit of a - unit digit of b

Unit digit of $a \times b$ = unit digit of $a \times$ unit digit of b

Ex: What are the unit digit of 435×433 .

Sol: Unit digit of $a \times b$ = unit digit of $a \times$ unit digit of b

$$5 \times 3 = 15 \text{ so, unit digit is } 5$$

Cyclicity

Cyclicity in the number system means the repeating pattern of digits or remainders when a number is raised to higher powers. The unit digit remains unchanged for all powers.

$$0 \rightarrow 0 \quad 1 \rightarrow 1$$

$$5 \rightarrow 5 \quad 6 \rightarrow 6$$

Cycle of Length 2-The unit digit alternates between two values.

$$4 \rightarrow 4, 6$$

When the power is odd, the unit digit is 4, and when the power is even, the unit digit is 6.

$$9 \rightarrow 9, 1$$

When the power is odd, the unit digit is 9, and when the power is even, the unit digit is 1.

Cycle of Length 4- The unit digit repeats after four powers.

$$2 \rightarrow 2, 4, 8, 6$$

$$3 \rightarrow 3, 9, 7, 1$$

$$7 \rightarrow 7, 9, 3, 1$$

$$8 \rightarrow 8, 4, 2, 6$$

$$\text{Let } N = x^y$$

To find the unit digit of (N) , we only need to consider the unit digit of the base number (x) .

The unit digit of an exponential expression can be determined by finding the remainder when the power is divided by 4.

Type 3: Cyclicity - Unit Digit Based on Arithmetic Equations



Ex: If $x = (164)^{169} + (333)^{337} - (727)^{726}$ then what is the unit digit of x ?

Sol: In the expression, the **first term** has an odd power of 4, so the unit digit of the first term is 4. For the **second term**, dividing 337 by 4 gives a remainder of 1 so the unit digit of the second term is 3. For the **third term**, dividing 726 by 4 gives a remainder of 2; hence, the unit digit of the third term is 9.

So, unit digit of the expression $4 + 3 - 9 = -2$. If the unit digit comes out negative, add 10 to obtain the correct unit digit. Unit digit is $10 - 2 = 8$.

Ex: What is the unit digit of $1^5 + 2^5 + 3^5 + 4^5 + \dots + 20^5$

Sol: In each term, the cyclicity is 1. Therefore, for every term, the units digit is the same as the number itself.

The Unit digit for 1 to 10 is zero.

$$= (1 + 2 + 2.. + 9 + 0)$$

$$+ (1 + 2 + 3.. + 9 + 0) = 0$$

Ex: Find the unit digit $x = 187^{280} \times 529^{320} \times 343^{236}$

Sol: If remainder is 0 then put power equal to the 4.

For the first terms - 7^4 cyclicity is 1

For the second terms - power of 9 is even so the unit digit is 1

For the third terms - 3^4 cyclicity is 1

Unit digit is $= 1 \times 1 \times 1 = 1$

Ex: What is the unit digit in the expansion of $(57242)^{9 \times 7 \times 5 \times 3 \times 1}$?

Sol: We only check only for digit 2. Power will be divided by 4.

$$= 2^{1 \times (-1) \times 1 \times (-1) \times 1} = 2^1$$

So, the unit digit is 2

Type 4: Counting based Questions (a digit or page counting or key strokes)



1 to 9 $\rightarrow$ required digit is = 9

10 to 99 $\rightarrow 90 \times 2 = 180$

Ex: What is the number of digits required for numbering a book with 428 pages?

Sol: 1 to 9 $\rightarrow$ required digit is = 9

10 to 99 $\rightarrow 90 \times 2 = 180$

100 to 428 $= (428 - 100 + 1) = 329 \rightarrow 329 \times 3 = 987$

Total number of digits required = $9 + 180 + 987 = 1176$

Type 5: Perfect Square Based Questions



How to check if a number is a perfect square (it shows the possibility)

1. The last two digits of any perfect square number must be among the squares from 1 to 24.
2. The units digit should not be 2, 3, 7, or 8.
3. The number of zeros in the number and the denominator should both be even.
4. After dividing a perfect square by 9, the remainder must be 0, 1, 4, or 7.

Ex: Is it possible that 562576 is a perfect square or not?

Sol: The number ends in 76, which is possible for a perfect square. The unit digit is 6, so it is possible for the number to be a perfect square.

Sum of digit of 562576 = 21

after dividing remainder is 3.

So, this number is not perfect.

Type 6 Decimal to fraction conversion



- The number of zeros in the denominator is equal to the number of digits after the decimal point.

$$0.abc = \frac{abc}{1000}$$

- The number of 9 in the denominator is equal to the number of digits after the decimal point.

$$0.\overline{abc} = \frac{abc}{999}$$

- When some digits are not covered by the overline

$$0.a\overline{bc} = \frac{abc - a}{990}$$

$$0.ab\overline{cd} = \frac{abcd - ab}{9900}$$

- Mix concept

$$a.b\overline{cd} = a + \frac{bcd - b}{990} = \frac{abcd - ab}{990}$$

Ex: If $A = 0.3\overline{12}$, $B = 0.4\overline{15}$ and $C = 0.30\overline{9}$ what is the value of $A + B + C$.

Sol:

$$A + B + C = \frac{312 - 3}{990} + \frac{415 - 4}{990} + \frac{309 - 30}{900}$$

$$A + B + C = \frac{720}{999} + \frac{279}{900}$$

$$A + B + C = \frac{10269}{9900} = \frac{1141}{1100}$$



Type 7: Number of zeros

- Zeros at the end of a number are determined by the number of 10s and in its factorization, primarily based on the pairs of 5s and 2s.
- **Factorial** is a mathematical operation defined for non-negative integers.

- For a positive integer n , the factorial of n , denoted by $n!$, is the product of all positive integers from 1 to n .

$$n! = n \times (n-1) \times (n-2) \times \dots \times 1$$

$$0! = 1, 1! = 1$$

- ✓ The unit digit of numbers after $4!$ is zero.
 - ✓ $4!$ and all factorials after it are divisible by 4.
 - ✓ Product of ' n ' consecutive natural number is divisible by n
- **Power of number contained in a factorial:** Highest power of a prime number ' p ' contained in $n!$ is given by
- $$= \left[\frac{n}{p} \right] + \left[\frac{n}{p^2} \right] + \left[\frac{n}{p^3} \right] + \dots$$
- The product of n consecutive natural number is always divisible by $n!$

Ex: The three numbers 24, 25, and 26 are divisible by.

Sol: The product of n consecutive natural number is always divisible by $n!$

Means 24, 25, 26 is divisible by 3!

Ex: Find the number of trailing zeros in 100!

Sol: In $100!$ factors of 2 are abundant, we only count factors of 5.

Each multiple of 5 contributes at least one factor of 5. Numbers like 25, 50, 75, 100 contain an extra factor of 5 because $25 = 5^2$.

$$\left[\frac{100}{5} \right] + \left[\frac{100}{25} \right] + \left[\frac{100}{125} \right] = 20 + 4 + 0 = 24$$

Number of trailing zeros = 24

Ex: Find the number of trailing zeros in $2 \times 4 \times 6 \dots \times 250$.

Sol: factors of 2 are abundant, we only count factors of 5.

$$2 \times 4 \times 6 \dots \times 250$$

$$= (2 \times 1) \times (2 \times 2) \dots (2 \times 125)$$

$$(2 \times 1) \times (2 \times 2) \dots (2 \times 125)$$

$$= 2^{125} (1 \times 2 \times \dots \times 125)$$

$$2^{125} (1 \times 2 \times \dots \times 125) = 2^{125} \times 125!$$

$$\left[\frac{125}{5} \right] + \left[\frac{125}{25} \right] + \left[\frac{125}{125} \right] = 25 + 5 + 1 = 31$$

Number of trailing zeros = 31

Divisibility

Number	Divisibility Rule
2	Last digit is 0, 2, 4, 6, 8
3	Sum of digits divisible by 3
4	Last two digits form a number divisible by 4
5	Last digit is 0 or 5
25	Last two digit is 00 or divisibly by 25
6	Number divisible by both 2 and 3
7	Subtract twice the last digit from the remaining; result divisible by 7
8	Last three digits divisible by 8
9	Sum of digits divisible by 9
11	Difference between sum of digits at even and odd places is 0 or divisible by 11

Special cases

$$1. \quad 1001 = 7 \times 11 \times 13$$

$$1001 \times abc = abcabc$$

$$2. \quad 10101 = 3 \times 7 \times 13 \times 37$$

$$10101 \times ab = ababab$$

Type 8: Based on

Divisibility rules

Ex: A number N is formed by writing 9 for 99 times. What is the remainder if N is divided by 13?

Sol: When any number is repeated n times, the combination of the 6-digit number is divisible by 7, 11, and 13.

96 times 9 will be divisible by 13, and only three 9s will remain.

$$\frac{999}{13} \Rightarrow R \rightarrow 11$$

Ex: Find the greatest possible value of $(a + b)$ for which the 8-digit number 143b203a is divisible by 15.

Sol: Divisibility of 3 - if sum of its digits is divisible by 3.

Divisibility of 5 - if the last number is either 0 or 5, the entire no. is divisible by 5.

Factors of 15 = (3×5) Hence, the number should be divisible by both 3 and 5.

So, a can be 0 or 5. But since the question asks for greatest value, a must be 5.

After that sum of the number is = $18 + b$

For greatest = $b = 9$

$$\text{So, } (a + b) = (9 + 5) = 14$$



Ex: If the 9-digit number $72x8431y4$ is divisible by 36, find the value of $\left(\frac{x}{y} - \frac{y}{x}\right)$ for the smallest possible value of y , where x and y are natural numbers.

Sol: The divisibility rule of 36 is that the number must be divisible by 4 and 9.

Last two-digit number = $y4$, on putting $y = 2$, the last two-digit number becomes 24, Therefore, the value of $y = 2$. (y is the smallest possible value)

For divisibility rule of 9,
sum of number = $31 + x$

So, $x = 5$

Now the value

$$\left(\frac{x}{y} - \frac{y}{x}\right) = \left(\frac{5}{2} - \frac{2}{5}\right) = 2\frac{1}{10}$$



Type 9: Divisibility of expressions

Expression	'n' is odd	'n' is even
$x^n - y^n$	$(x - y)$	$(x + y)(x - y)$
$x^n + y^n$	$(x + y)$	Can't say

Ex: What is the remainder if $(17^{26} - 11^{26})$ is divided by 42?

Sol: n is even. So, $(x + y)(x - y)$

$$= \frac{(17 + 11)(17 - 11)}{42} = \frac{28 \times 6}{42},$$

Remainder = 0

Key Point: If ' n ' is odd and $a, b, c \dots z$ are consecutive natural number then $(a^n + b^n + \dots + z^n)$ is divisible by $(a + b + c \dots z)$

Ex: $11^5 + 12^5 + 13^5$ is divisible by?

Sol- using concept

$$11 + 12 + 13 = 36$$

So, the expression is divisible by 36

Type 10: Digit and its Reverse Form 2-digit and 3 digit and its reverse form



1. For 2-digit

Let, two-digit original number = $10x + y$, Reversed number = $10y + x$

Sum of original and reserved number
 $(10x + y) + (10y + x) = 11(x + y)$

Difference of original and reversed number
 $(10x + y) - (10y + x) = 9(x - y)$

2. For 3-digit

Let, Hundred digit = x , Tenth digit = y , Unit digit = z So,

original number = $100x + 10y + z$

Hundred and unit digits interchange,

new number = $100z + 10y + x$

Difference of original and reverse number = $99(x - z)$

Ex: The sum of the two-digit number and the number obtained by inter-changing the digit is 99. If the difference of digits is 1, then the number is;

Sol: Let, number = $10x + y$

$$(10x + y) + (10y + x) = 99$$

$$x + y = 9$$

$$x - y = 1$$

After the equation $x = 5, y = 4$

So, the number is

$$= 10 \times 5 + 4 = 54$$

Ex: What will be the smallest natural number to be filled in the blank for the number $23x45678$ to be divisible by 22.

Sol: The divisibility rule for 22 is based on divisibility by both 2 and 11. A number is divisible by 11 if the difference between the sum of its digits in odd positions and the sum of its digits in even positions is divisible by 11.

Odd positions sum: $3 + 4 + 6 + 8 = 21$

Even positions sum: $2 + x + 5 + 7 = 14 + x$

$$21 - (14 + x) = 7 - x$$

$$7 - x = 11k, \text{ for some integer } k$$

For the smallest x , let's try $k = 0$

$$7 - x = 0 \Rightarrow x = 7$$

Thus, the smallest value of x that makes the number $23x45678$ divisible by 22 is 5.

Ex: How many numbers are there from 700 to 950 (including both) which are neither divisible by 3 nor by 7?

Sol:

Neither divisible by 3 nor by 7

$$= \text{Total} - (\text{Divisible by 3 or 7}) = \text{Total} - [N(3) + N(7) - N(21)]$$

Total number from 700 to 950 = 251

$$\text{Divisible by 3} = \frac{251}{3} \approx 83$$

$$\text{Divisible by 7} = \frac{251}{7} \approx 35$$

$$\text{Divisible by 21} = \frac{251}{21} \approx 11$$

Required number

$$= 251 - (83 + 35 - 11) = 144$$

Factorization

A factor of a number is a whole number that can be multiplied by another whole number to result in the original number. In other words, a factor divides the number evenly without leaving any remainder.

$$\text{Let } N = a^p \times b^q \times c^r$$

$$\text{Total number of factors} = (p + 1)(q + 1)(r + 1)$$

$$\text{Total number of even factors} = p(q + 1)(r + 1)$$

$$\text{Total number of odd factors} = (q + 1)(r + 1)$$

$$\text{Sum of all factors} = (a^0 + a^1 + \dots + a^p)(b^0 + b^1 + \dots + b^q)(c^0 + c^1 + \dots + c^r)$$

$$\text{Sum of even factors} = (2^1 + 2^2 + \dots + 2^p)(b^0 + b^1 + \dots + b^q)(c^0 + c^1 + \dots + c^r)$$

$$\text{Sum of odd factors} = 2^0(b^1 + b^2 + \dots + b^q)(c^1 + c^2 + \dots + c^r)$$

$$\text{Average of factor} = \frac{\text{sum of factor}}{\text{number of factor}}$$

$$\text{Sum of reciprocal of factors} = \frac{\text{sum of factor}}{\text{given number}}$$

$$\text{Number of prime factors} = p + q + r$$

$$\text{Number of distinct prime factor} = \text{no of prime number in factor}$$

$$\text{Number of composite numbers} = \text{total} - \text{distinct prime number} - 1$$

- Every number has positive and negative factors.
- All numbers in the form $(\text{prime})^2$ have 3 factors.

Type 11: Find the Number of Factors Based



Ex: Find all types of factors and the sum of all types of factors for $N = 3600$.

$$\text{Sol: } N = 3600 = 2^4 \times 3^2 \times 5^2$$

- Total number of factors $= 5 \times 3 \times 3 = 45$
- Number of even factors
 $2 \times (1800) = 2(2^3 \times 3^2 \times 5^2)$
 $= 4 \times 3 \times 3 = 36$
- Number of odd factors $= 3 \times 3 = 9$
- Number of prime factors $= 4 + 2 + 2 = 8$
- Number of distinct Prime factor (visible in factorization) $= 1 + 1 + 1 = 3$
- Number of Composite numbers
 $= 45 - 3 - 1 = 41$
- Perfect square $= (2^2)^2(3^2)^1(5^2)^1$
 $= 3 \times 2 \times 2 = 12$

- Perfect cube $= (2^3)^1 = (1 + 1) = 2$
- Sum of all factor
 $= (2^0 + 2^1 + 2^2 + 2^3 + 2^4)(3^0 + 3^1 + 3^2)(5^0 + 5^1 + 5^2)$
 $= 12493$
- Sum of even factor
 $= (2^1 + 2^2 + 2^3 + 2^4)(3^0 + 3^1 + 3^2)(5^0 + 5^1 + 5^2) = 12090$
- Sum of odd factor
 $= (3^0 + 3^1 + 3^2)(5^0 + 5^1 + 5^2) = 403$
- Sum of prime $= 2 + 3 + 5 = 10$
- Sum of composite number
 $= \text{total sum} - (\text{sum of prime} + 1) = 12493 - 10 - 1 = 12482$
- Perfect square
 $= (2^0 + 2^2 + 2^4)(3^0 + 3^2)(5^0 + 5^1) = 5460$

Ex: Calculate the total number of prime factors in the expression $(4^{11} \times 5^5 \times 3^2 \times 13^2)$

Sol:

$$(4^{11} \times 5^5 \times 3^2 \times 13^2) = 2^{22} \times 3^2 \times 5^5 \times 13^2$$

Total number of prime factors

$$= 22 + 2 + 5 + 2 = 31$$

Ex: How many factors of $(30^{16} \times 16^{18} \times 20^{21})$ are perfect square as well as perfect cube?

$$\text{Sol: } 30^{16} \times 16^{18} \times 20^{21} = 2^{130} \times 3^{16} \times 5^{37}$$

When asked to check for a perfect square or perfect cube, check if the power is a multiple of 6.

$$2^{130} \times 3^{16} \times 5^{37}$$

$$= 2^{126} \times 2^4 \times 3^{12} \times 3^4 \times 5^{36} \times 5^1$$

$$2^{126} \times 2^4 \times 3^{12} \times 3^4 \times 5^{36} \times 5^1$$

$$= (2^6)^{21} \times 2^4 \times (3^6)^2 \times 3^4$$

$$\times (5^6)^6 \times 5^1$$

The total number of factors that are both perfect squares and perfect cubes

$$= (21 + 1)(2 + 1)(6 + 1) = 462$$

Ex: How many factors of the number $2^8 \times 3^6 \times 5^4 \times 10^5$ are multiple of 120.

$$\text{Sol: } N = 2^8 \times 3^6 \times 5^4 \times 10^5 = 2^{13} \times 3^6 \times 5^9$$

$$\frac{N}{120} = \frac{2^{13} \times 3^6 \times 5^9}{2^3 \times 3^1 \times 5^1} = 2^{10} \times 3^5 \times 5^8$$

$$\text{Number of factors} = 11 \times 6 \times 9 = 594$$

Remainder

Let suppose number N when divided by divisor D , leaves the remainder as R and quotient as Q .

$$N = D \times Q + R$$

Dividend = Divisor $\times$ quotient + Remainder

If a gives r remainder when divided by n , then ka gives kr remainder when divided by n .

If a and b give r_1 and r_2 remainders respectively when divided by n then.

$a + b$ divided by n , gives remainder $r_1 + r_2$

$a \times b$ divided by n , gives remainder $r_1 \times r_2$

Type 12: Question based on remainder theorem

Ex: A number being divided by 52 gives a remainder 45. If the number is divided by 13, the remainder will be?

Sol: Since 13 is a multiple of 52, we can directly divide the remainder of one by the divisor.

$$\frac{45}{13} \Rightarrow R \rightarrow 6$$

Ex: A number when divided by 12 leaves 5 remainder. What is the remainder if square of this number is divided by 8.

Sol: When a mathematical operation is performed on a number, the same operation can also be performed on the remainder.

$$\frac{5^2}{8} = \frac{25}{8} \Rightarrow R \rightarrow 1$$

Ex: If the dividend is 45, the divisor is 8, and the quotient is 5, find the remainder.

$$\text{Sol- } N = D \times Q + R$$

$$45 = 8 \times 5 + \text{Remainder}$$

$$\text{Remainder} = 45 - 40 = 5$$

Type 13: Important concept of remainder

$$\frac{(x+a)^n}{x} \Rightarrow R \rightarrow a^n$$

$$\frac{(x+1)^n}{x} \Rightarrow R \rightarrow 1$$

$$\frac{(x-1)^n}{x} \Rightarrow R \rightarrow (1)^n$$

$$R = 1 \text{ when } n \text{ is even}$$

$$R = -1 \text{ when } n \text{ is odd}$$

Special case:

$$\frac{4}{6} \Rightarrow R \rightarrow 4$$

$$\frac{4^n}{6} \Rightarrow R \rightarrow 4$$

Ex: What is the remainder when 2^{75} is divided by 14?

Sol:

$$\frac{2^{75}}{15} = \frac{(2^4)^{18} \cdot 2^3}{15} = \frac{(16-1)^{18} \times 8}{15} \Rightarrow R \rightarrow 8$$

Ex: What is the remainder if 2^{192} is divided by 6?

Sol: Using the concept

$$\frac{2^{192}}{6} = \frac{4^{96}}{6} \Rightarrow R \rightarrow 4$$

Ex: What is the remainder when 37^{47} is divided by 19.

Sol:

$$\frac{37^{47}}{19} = \frac{(38-1)^{47}}{19} \Rightarrow R \rightarrow -1$$

Remainder cannot be negative, so the remainder is subtracted from the divisor.

$$R \rightarrow -1 \Rightarrow R \rightarrow 19 - 1 = 18$$

Type 14: Fermat's little theorem

$$\text{If } \frac{a^{p-1}}{p}, \Rightarrow R \rightarrow 1$$

Conditions

1. P is prime number.
2. a, P are co-prime.

Ex: What is the remainder when 21^{47} is divided by 47?

Sol: 21 and 47 are coprime number so we can use Fermat's little theorem.

$$\frac{21^{47}}{47} = \frac{21^{46} \times 21}{47} \Rightarrow R \rightarrow 21$$

Type 15: Euler's (totient) theorem

$$\frac{A^X}{d} \Rightarrow R \rightarrow 1$$

Where A and d are co-prime.

X – number of numbers less than d and co-prime to d .

$$X = d \left(1 - \frac{1}{p^1}\right) \left(1 - \frac{1}{p^2}\right) \dots \dots \dots$$

Where P^1 and P^2 are distinct prime factors of d .

Ex: If ϕ is the Euler's totient functions, then $\phi(92)$ is.

Sol: $92 = 2^2 \times 23$

$$\phi(92) = 92 \left(1 - \frac{1}{2}\right) \times \left(1 - \frac{1}{23}\right) = 44$$

Ex: 7^{82} is divided by 11, R=?

Sol: To find the remainder when 7^{82} is divided by 11, we can use Fermat's Little Theorem, which states that if p is a prime number and a is an integer not divisible by p , then

$$\phi(11) = 11 \left(1 - \frac{1}{11}\right) = 10$$

82 is divided by 10 , and the remainder is

$$2 \cdot \frac{7^2}{11} \Rightarrow R \rightarrow 5$$

Type 16: Wilson's theorem



If P is prime number, then,

$$\frac{(P-1)!}{P} \Rightarrow R \rightarrow P-1$$

$$\frac{(P-2)!}{P} \Rightarrow R \rightarrow 1$$

Ex: What will be the remainder when $568!$ is divided by 569 .

Sol- Using the Wilson theorem

$$\frac{568!}{596} = \frac{(569-1)!}{P} \Rightarrow R \rightarrow 569-1$$
$$= 568$$



Toppersnotes
Unleash the topper in you



CHAPTER

LCM- least common multiple

- The smallest number that is exactly divisible by each of the given numbers is called their LCM.

Example:

Factor of 12 - 12, 24, 36, 48, ...

Factor of 16 - 16, 32, 48, 64, ...

Common multiple - 48, 96...

Least common multiple - 48

HCF- highest common factor

- The largest number that exactly divides each of the given numbers is called their HCF.

Example

Factors of 24 - 1, 2, 3, 4, 6, 8, 12, 24

Factors of 36 - 1, 2, 3, 4, 6, 9, 12, 18, 36

Common factors of 24 and 36 - 1, 2, 3, 4, 6, 12

Highest common factor - 12

Important Exam - Oriented Types**Type 1: Least common Multiple - based**

Find the smallest number that is exactly divisible by x, y and z .

$= \text{LCM of } (x, y, z)$

1. Find the smallest number that is divided by x, y and z and leaves remainder r in each case.

$= \text{LCM of } (x, y, z) + r$

2. Find the smallest number that is divisible by x, y , and z , and leaves remainders of a, b , and c , respectively.

$= \text{LCM of } (x, y, z) - k$

Where $k = (x - a) = (y - b) = (z - c)$

Ex: Find the smallest number which when divided 5, 6, 7 and 8 gives remainder 3 in each case.

Sol: Smallest number $= \text{LCM of } (x, y, z) + r$
 $= \text{LCM}(5, 6, 7, 8) + 3 = 840 + 3 = 843$

Ex: Find the smallest number that, when divided by 25, 15, and 30, leaves remainders of 21, 11, and 26, respectively.

Sol:

$$k = (25 - 21) = (15 - 11) = (30 - 26) = 4$$

$$= \text{LCM of } (25, 15, 30) - 4 = 146$$

Ex: When the numbers 12, 16, 18, 20, and 25 divide the least number x , the remainder in each case is 4, but x is divisible by 7. What is the digit at the thousands' place in x

Sol:

$$\text{LCM} = \text{LCM of } (12, 16, 18, 20, 25) = 3600$$

Given, 3600 should be divisible by 7.

Let $(3600k + 4)$ is divisible by 7.

So, we can write

$$\frac{3600k + 4}{7}$$

Put $k = 1, 2, 3, 4, 5, \dots$ if we put $k = 5$ then expression is divisible by 7.

$$\text{Number} = 3600 \times 5 + 4 = 18004$$

Digit at thousands' place in 18004 is 8.

Type 2: Highest Common factor - Based

1. Find the largest number which is divisible by x, y and z exactly.

$= \text{HCF of } (x, y, z)$

2. Largest number which can divide x, y and z and leaves remainder r .

$= \text{HCF of } (x, y, z) + r$

If r is absent.

$= \text{HCF of } |x - y|, |y - z|, |z - x|$

3. Largest number which can be divide x, y, z and leaves remainder a, b, c respectively.

$= \text{HCF of } (x - a)(y - b)(z - c)$

Ex: If $X = 2^3 \times 3^{10} \times 5^1$ and $Y = 2^5 \times 3^1 \times 7^1$, then find the HCF of X and Y ?

Sol: The largest number that exactly divides each of the given numbers is called their HCF.

$$\text{HCF} = 2^3 \times 3^1$$

Ex: Find the greatest possible length which can be used to measure exactly the length 7m, 3m 85cm, 12m 95cm.

Sol:

$$= \text{HCF of } (700, 385, 1295)$$

$$= \text{HCF of } (x - a)(y - b)(z - c)$$

$$\text{HCF of } (700 - 385)(1295 - 385)(1295 - 700)$$

$$= \text{HCF of } (315)(1225)(595)$$

$$\text{HCF of } (315)(1225)(595) = 35 \text{ cm}$$

Type 3: LCM and HCF of fractions



$$\text{LCM of } \left(\frac{a}{b}, \frac{c}{d}, \frac{e}{f}\right) = \frac{\text{LCM of } (a, c, e)}{\text{HCF of } (b, d, f)}$$

$$\text{HCF of } \left(\frac{a}{b}, \frac{c}{d}, \frac{e}{f}\right) = \frac{\text{HCF of } (a, c, e)}{\text{LCM of } (b, d, f)}$$

Key point - The fraction must be in its simplest form.

Ex: What is the LCM of 3.6, 1.8 and 0.144?

Sol:

$$\frac{36}{10} = \frac{18}{5}, \quad \frac{18}{10} = \frac{9}{5}, \quad \frac{144}{1000} = \frac{18}{125}$$

$$\text{LCM of } \left(\frac{18}{5}, \frac{9}{5}, \frac{18}{125}\right) = \frac{\text{LCM of } (18, 9, 18)}{\text{HCF of } (5, 5, 125)}$$

$$= \frac{18}{5} = 3.6$$

LCM & HCF of powers

$$a^p + 1, \quad a^q + 1$$

$$\text{LCM of } (a^p + 1)(a^q + 1) = a^{\text{LCM of } (p, q)} + 1$$

$$\text{HCF of } (a^p + 1)(a^q + 1) = a^{\text{HCF of } (p, q)} + 1$$

Ex: What is the HCF $(2^{36} - 1)$ and $(2^{45} - 1)$.

Sol:

$$\text{HCF of } (a^p + 1)(a^q + 1) = a^{\text{HCF of } (p, q)} + 1$$

$$\text{HCF of } (2^{36} - 1)(2^{45} - 1)$$

$$= 2^{\text{HCF of } (36, 45)} + 1 = 2^9 + 1 = 511$$

Type 4: Relationship between LCM and HCF Based



Let the 2 numbers are

$$N_1 = ha, N_2 = hb \rightarrow 'a' \text{ and } 'b' \text{ are coprime} \\ \rightarrow h(a, b) = 1$$

$$\text{LCM} \times \text{HCF} = N_1 \times N_2$$

$$\text{LCM} = hab$$

$$\text{sum} = (a + b)h$$

$$\text{difference} = (a - b)h$$

$$\text{Product} = h^2 ab$$

$$\frac{\text{LCM}}{\text{HCF}} = \frac{ab}{1}$$

Ex: Two numbers are in the ratio of 5:7. the product of their LCM and HCF is 12635. Then the sum of the number will be.

Sol: Let HCF = x , then numbers are $5x, 7x$

$$5x \times 7x = 12635$$

$$x = 19$$

$$\text{sum} = (a + b)h = (5 + 7)19 = 228$$

Ex: The sum of and difference between the LCM and HCF of two numbers are 512 and 496, respectively. If one number is 72, then the other number is.

Sol: Let second number is N_1

$$\text{LCM} + \text{HCF} = 512$$

$$\text{LCM} - \text{HCF} = 496$$

After solving the equations

$$\text{LCM} = 504, \quad \text{HCF} = 8$$

$$72 \times N_2 = 504 \times 8$$

$$N_2 = 56$$

Type 5: LCM and HCF questions based on ratios



Ex: Three numbers are in the ratio $\frac{1}{2} : \frac{2}{3} : \frac{3}{4}$. The difference between the largest and the smallest number is 21. The sum of all the numbers is.

Sol:

$$\text{largest number} - \text{smallest number} = 21$$

$$\frac{1}{2} : \frac{2}{3} : \frac{3}{4} = 6 : 8 : 9$$

Now, let the ratio be x then, numbers are $6x, 8x$ and $9x$

$$9x - 6x = 21$$

$$x = 7$$

Sum of all three numbers

$$= (6 + 8 + 9) \times 7 = 161$$

Type 6: Questions based on bells and traffic signals



Ex: There are three traffic signals.

Each signal changes colour from green to red and then from red to green. The first signal takes 25 seconds, the second signal takes 39 seconds and the third signal takes 60 seconds to change the colour from green to red. The durations for green and red colours are same. At 2:00 p.m., they together turn green. At what time will they change to green next, simultaneously?

Sol: The three traffic signals turn from green to red at an interval of 25 second, 39 second, and 60 seconds.

$$LCM(25,39,60) = 3900 \text{ sec} = 65 \text{ min}$$

$$\text{Total} = 2 \text{ times} = 65 + 65 = 130 \text{ min}$$

Time = 4:10 P.M.

Ex: Four ropes of lengths 102m, 119m, 153m and 204 m are to be cut into parts of equal length. Each part must be as long as possible. What is the maximum number of pieces that can be cut?

Sol:

HCF = equal length in which ropes can be cut

$$HCF(102, 119, 153, 204) = 17$$

$$\begin{aligned} \text{Total parts} &= \frac{102}{17} + \frac{119}{17} + \frac{153}{17} + \frac{204}{17} \\ &= 6 + 7 + 9 + 12 = 34 \end{aligned}$$

Type 7: LCM and HCF of polynomials



Ex: What is the HCF of $(x^8 - y^8)$ and $(x^7 - y^7 + x^5y^2 - x^2y^5)$.

Sol:

$$\begin{aligned} (x^8 - y^8) &= (x^4 + y^4)(x^2 + y^2)(x + y)(x - y) \\ (x^7 - y^7 + x^5y^2 - x^2y^5) & \end{aligned}$$

$$= x^7 + x^5y^2 - y^7 - x^2y^5$$

$$x^7 + x^5y^2 - y^7 - x^2y^5 = (x^2 + y^2)(x^5 - y^5)$$

$$(x^2 + y^2)(x^5 - y^5)$$

$$= (x^2 + y^2)(x - y)(x^4 + x^3y$$

$$+ x^2y^2 + xy^3 + y^4)$$

$$\text{HCF of } (x^8 - y^8)(x^7 - y^7 + x^5y^2 - x^2y^5)$$

$$= (x^2 + y^2)(x - y)$$

$$\text{HCF} = (x^2 + y^2)(x - y)$$

$$= x^3 - x^2y + xy^2 - y^3$$



Toppersnotes
Unleash the topper in you

3

CHAPTER

Surds and Indices

Surds



A surd is an irrational number expressed in the form of a root ($\sqrt[n]{}$) that cannot be simplified into a rational number or cannot be expressed in rational form ($\frac{p}{q}$).

$\sqrt{2}, \sqrt{3}, \sqrt{5} \rightarrow$ surds

$\sqrt{4} = 2$ (not a surd, because it is rational)

Properties:

- $\sqrt[n]{a} = a^{\frac{1}{n}}$
- $\sqrt[n]{ab} = \sqrt[n]{a} \times \sqrt[n]{b} = a^{\frac{1}{n}} \times b^{\frac{1}{n}} = (ab)^{\frac{1}{n}}$
- $\sqrt[n]{\frac{a}{b}} = \frac{\sqrt[n]{a}}{\sqrt[n]{b}} = \left(\frac{a}{b}\right)^{\frac{1}{n}}$
- $(\sqrt[n]{a})^n = a^{\frac{n}{n}} = a$
- $(\sqrt[n]{a})^m = a^{\frac{m}{n}} = \sqrt[n]{a^m}$
- $\sqrt[n]{\sqrt[m]{a}} = \sqrt[n]{a^{\frac{1}{m}}} = a^{\frac{1}{mn}} = \sqrt[nm]{a}$
- $\sqrt{2} = 1.414$
- $\sqrt{3} = 1.732$
- $\sqrt{5} = 2.236$

Indices



An index (or exponent) is a number that shows how many times a base number is multiplied by itself. It represents repeated multiplication in a compact form. $a, a, a, \dots$ to m terms is written as a^m . Where a is base and m is index.

Properties:



- $a^m \times a^n = a^{m+n}$
- $(a^m)^n = a^{mn}$
- $a^m \div a^n = a^{m-n}$
- $\frac{1}{a^{-n}} = a^n$
- $(ab)^n = a^n b^n$
- $a^0 = 1$, if $a \neq 0$
- $a^1 = a$
- $a^x = a^y \rightarrow x = y$

Rationalization:



Rationalization is the process of removing an irrational number (like $\sqrt{2}, \sqrt{3}$) from the denominator of a fraction by multiplying the numerator and denominator by a suitable irrational expression.

$$= \frac{1}{\sqrt{3} + 1}$$

Multiply by the conjugate ($\sqrt{3} - 1$)

$$= \frac{1}{\sqrt{3} + 1} \times \frac{\sqrt{3} - 1}{\sqrt{3} - 1} = \frac{\sqrt{3} - 1}{2}$$

Type 1: Question based on basic identities

Ex: If $\frac{\sqrt{6}-2}{\sqrt{6}+2} = a\sqrt{6} + b$, find the value of $b^2 + a^2$?

Sol: Rationalize the denominator

$$\frac{\sqrt{6}-2}{\sqrt{6}+2} \times \frac{\sqrt{6}-2}{\sqrt{6}-2} = 5 - 2\sqrt{6}$$

$$a = -2, \quad b = 5$$

$$b^2 + a^2 = 25 + 4 = 29$$

Ex: Find the value of $\left[\left[\frac{-1^2}{2}\right]^{-2}\right]^{-1}$

Sol: Using this property, the powers are multiplied. $(a^m)^n = a^{mn}$

$$\left[\left[\frac{-1^2}{2}\right]^{-2}\right]^{-1} = \left[\frac{-1}{2}\right]^4 = \frac{1}{16}$$

Ex: Find the value of

$$\frac{2}{\sqrt{7} + \sqrt{5}} + \frac{7}{\sqrt{12} - \sqrt{5}} - \frac{5}{\sqrt{12} - \sqrt{7}}$$

Sol: Rationalization of each term

$$\frac{2}{2} [\sqrt{7} - \sqrt{5} + \sqrt{12} + \sqrt{5} - \sqrt{12} + \sqrt{7}] = 0$$

Ex: Find A×B

$$A = \frac{\sqrt{0.0004} \times \sqrt[3]{0.000008}}{\sqrt[4]{16000} \times \sqrt[3]{125000} \times \sqrt[4]{810}}$$

$$B = \frac{\sqrt[3]{0.729} \times \sqrt[4]{0.0016}}{\sqrt{0.16}}$$

Sol:

$$A \times B = \frac{0.02 \times 0.02}{2 \times 10 \times 50 \times 3} \times \frac{0.9 \times 0.2}{0.4}$$

$$A \times B = 6 \times 10^{-8}$$

Ex: Arranging the following in ascending order $3^{34}, 2^{51}, 7^{17}$.

Sol: $3^{34} = (3^2)^{17} = 9^{17}$

$2^{51} = (2^3)^{17} = 8^{17}$

Since for the same positive exponent (17):

$7 < 8 < 9$

$7^{17} < 8^{17} < 9^{17}$

Replace back

$3^{34} > 2^{51} > 7^{17}$

Type 2: Infinite series

Multiplication and division

$$\sqrt[n]{a \sqrt[n]{a \sqrt[n]{a} \dots \dots \dots n \text{ times}}} = a^{(1-\frac{1}{2^n})},$$

$n = \text{number of roots}$

$$\sqrt[n]{a \sqrt[n]{a \sqrt[n]{a} \dots \dots \dots \text{up to } \infty}} = a$$

$$\sqrt[n]{a \div \sqrt[n]{a \div \sqrt[n]{a} \dots \dots \dots \text{up to } \infty}} = \sqrt[n]{a}$$

$$\sqrt[m]{a^n \sqrt[n]{b^m \sqrt[a]{a} \dots \text{up to } \infty}} = {}^{mn-1}\sqrt{a^n b}$$

$$\sqrt[n]{a \sqrt[n]{b \sqrt[n]{a \sqrt[n]{b} \dots \dots \dots \text{up to } \infty}}} = (a^2 b)^{\frac{1}{3}}$$

Addition and subtraction

General form

$$x = \sqrt[n]{a + b \sqrt[n]{a + b} \dots \dots \text{up to } \infty} = \frac{\sqrt{4a+b^2}+b}{2}$$

$$y = \sqrt[n]{a - b \sqrt[n]{a - b} \dots \dots \text{up to } \infty} = \frac{\sqrt{4a+b^2}-b}{2}$$

$x - y = b, \quad xy = a$

If $b = 1$

$$x = \sqrt[n]{a + \sqrt[n]{a + \dots \text{up to } \infty}} = \frac{\sqrt{4a+1}+1}{2}$$

$$y = \sqrt[n]{a - \sqrt[n]{a - \dots \text{up to } \infty}} = \frac{\sqrt{4a+1}-1}{2}$$

$x - y = 1, \quad xy = a, \quad x + y = \sqrt{4a+1}$

General form

$$x = \sqrt[n]{a + b \sqrt[n]{a - b} \dots \dots \text{up to } \infty}$$

$$= \frac{\sqrt{4a-3b^2}+b}{2}$$

$$y = \sqrt[n]{a - b \sqrt[n]{a + b} \dots \dots \text{up to } \infty}$$

$$= \frac{\sqrt{4a-3b^2}-b}{2}$$

If $b = 1$

$$x = \sqrt[n]{a + \sqrt[n]{a - \dots \text{up to } \infty}} = \frac{\sqrt{4a-3}+1}{2}$$

$$y = \sqrt[n]{a - \sqrt[n]{a + \dots \text{up to } \infty}} = \frac{\sqrt{4a-3}-1}{2}$$

$x - y = 1, \quad xy = a - 1,$

$x + y = \sqrt{4a-3}$

Ex: If $x = \sqrt{4\sqrt{4\sqrt{4} \dots \dots}}$, then find the value of x ?

Sol: using the concept

$$\sqrt[n]{a \sqrt[n]{a \sqrt[n]{a} \dots \dots \dots \text{up to } \infty}} = a$$

$x = 4$

Ex: Find $\sqrt{12\sqrt{12\sqrt{12} \dots \dots}}$?

Sol: In this case $n = 3$

$a^{(1-\frac{1}{2^3})} = 12^{\frac{7}{8}}$

Ex: Find the value of following

$$x = \sqrt{12 + \sqrt{12} \dots \dots \infty}$$

Sol: using the formula

$$x = \sqrt[n]{a + \sqrt[n]{a + \dots \text{up to } \infty}} = \frac{\sqrt{4a+1}+1}{2}$$

$$x = \frac{\sqrt{4 \times 12 + 1} + 1}{2} = 4$$

Trick: Whenever this type of question comes, the number should be broken in such a way that the product of the two numbers is equal to the given number, and the difference between them is 1.

If the question has a plus sign, the larger number will be the answer, and if it has a minus sign, the smaller number will be the answer.

$$12 = 4 \times 3$$

$$\text{So, } x = 4$$

Ex: Find $\sqrt{35 + 2\sqrt{35 + 2\sqrt{35 \dots \infty}}} = ?$

Sol: $a = 35, b = 2$

$$\frac{\sqrt{4a + b^2} + b}{2} = \frac{\sqrt{4 \times 35 + 4} + 2}{2} = 7$$

Ex: Find the value of given expression

$$\sqrt{7 + 2\sqrt{7 - 2\sqrt{7 \dots \infty}}} = ?$$

Sol: $a = 7, b = 2$

$$\frac{\sqrt{4a - 3b^2} + b}{2} = \frac{\sqrt{28 - 12} + 2}{2} = 3$$

Type 3: Comparison of Surds (Greater/Lesser) Based



To find Smallest / Largest in irrational numbers

When power is in fraction – Take LCM

Ex: Arrange in descending order

$$\sqrt[3]{3}, \sqrt[4]{4}, \sqrt[6]{6}, \sqrt[12]{12}$$

Sol: LCM (3,4,6,12) = 12

Multiply by LCM to each power

$$3^4 = 81, \quad 4^3 = 64, \quad 6^2 = 36, \quad 12^1 = 12$$

$$\sqrt[3]{3} > \sqrt[4]{4} > \sqrt[6]{6} > \sqrt[12]{12}$$

When power is in integer- Take HCF

Ex: Arrange in descending order

$$7^{200}, 9^{150}, 6^{250}, 5^{300}$$

Sol: HCF(200,150,250,300) = 50

Divide by HCF to each power

$$7^4 = 2401, 9^3 = 729, 6^5 = 7776, \quad 5^6 = 15625$$

$$5^{300} > 6^{250} > 7^{200} > 9^{150}$$

When sum is same

When the sign is plus, the maximum product corresponds to the largest number.

When the sign is minus, the maximum product corresponds to the smallest number.

Ex: Arrange in descending order

$$(\sqrt{8} + \sqrt{5}), (\sqrt{7} + \sqrt{6}), (\sqrt{9} + \sqrt{4}), (\sqrt{10} + \sqrt{3})$$

Sol- When the sign is plus, the maximum product corresponds to the largest number.

$$\begin{array}{cccc} 42 & 40 & 36 & 30 \\ (\sqrt{7} + \sqrt{6}) & > & (\sqrt{8} + \sqrt{5}) & > & (\sqrt{9} + \sqrt{4}) & > & (\sqrt{10} + \sqrt{3}) \end{array}$$

Ex: Arrange in ascending order

$$(\sqrt{8} - \sqrt{5}), (\sqrt{7} - \sqrt{6}), (\sqrt{9} - \sqrt{4}), (\sqrt{10} - \sqrt{3})$$

Sol: When the sign is minus, the maximum product corresponds to the smallest number.

$$\begin{array}{cccc} 42 & 40 & 36 & 30 \\ (\sqrt{7} - \sqrt{6}) & < & (\sqrt{8} - \sqrt{5}) & < & (\sqrt{9} - \sqrt{4}) & < & (\sqrt{10} - \sqrt{3}) \end{array}$$

When difference is same

Digit smaller, Number largest

Ex: Arrange in descending order

$$(\sqrt{3} - \sqrt{2}), (\sqrt{4} - \sqrt{3}), (\sqrt{5} - \sqrt{4}), (\sqrt{2} - \sqrt{1})$$

Sol:

$$(\sqrt{2} - \sqrt{1}) > (\sqrt{3} - \sqrt{2}) > (\sqrt{4} - \sqrt{3}) > (\sqrt{5} - \sqrt{4})$$

If the product is same

Sum/difference maximum, number largest

Ex: Arrange in descending order

$$(\sqrt{60} + \sqrt{6}), (\sqrt{15} + \sqrt{24}), (\sqrt{20} + \sqrt{18}), (\sqrt{30} + \sqrt{12})$$

Sol:

$$\begin{array}{cccc} 66 & 42 & 39 & 38 \\ (\sqrt{60} + \sqrt{6}) & > & (\sqrt{30} + \sqrt{12}) & > & (\sqrt{15} + \sqrt{24}) & > & (\sqrt{20} + \sqrt{18}) \end{array}$$

Type 4: Questions Based

on Making a Perfect Square



Ex: Find the value of $\sqrt{11 + 2\sqrt{18}}$.

Sol: Express the given expression in square form .18 must be broken into two numbers whose sum is 11

$$\sqrt{11 + 2\sqrt{18}} = \sqrt{9 + 2 + 2\sqrt{9}\sqrt{2}}$$

$$\sqrt{9 + 2 + 2\sqrt{9}\sqrt{2}} = \sqrt{(3 + \sqrt{2})^2}$$

$$\sqrt{11 + 2\sqrt{18}} = 3 + \sqrt{2}$$

$$3 + \sqrt{2} = 3 + 1.414 = 4.414$$

4

CHAPTER

Simplification



Simplification means reducing an expression to its simplest and most compact form by applying standard mathematical rules.

Order to solve

—	Vinculum/Line/Bar bracket
B	Bracket
O	Of
D	Division
M	Multiplication
A	Addition
S	Subtraction

Types of brackets and solving order

()	Small bracket
{ }	Curly bracket
[]	Square bracket

Important Exam – Oriented Types

Type 1: VBODMAS based



questions

Ex: The value of

$$441 \div \left[270 \div \frac{3}{7} \text{ of } 35 + \left(17 \div \frac{1}{3} \right) - \left(8\frac{1}{2} - \frac{5}{2} \right) \right]$$

is:

Sol: Using concept of BODMAS

$$= 441 \div \left[270 \div \left(\frac{3}{7} \times 35 \right) + (17 \times 3) - (17/2 - 5/2) \right]$$

$$= 441 \div \left[270 \div (3 \times 5) + 51 - \left(\frac{12}{2} \right) \right]$$

$$= 441 \div [270 \div 15 + 51 - 6]$$

$$= 441 \div 63 = 7$$

Ex: Simplify the following expression.

$$3\frac{2}{3} - 5\frac{7}{9} \div \frac{1}{3} \times 1\frac{2}{13}$$

$$\frac{2}{3} \text{ of } 1\frac{1}{3} \div \frac{1}{3}$$

Sol: Using concept of BODMAS

$$= 3\frac{2}{3} - 5\frac{7}{9} \div \frac{1}{3} \times 1\frac{2}{13}$$

$$= \frac{2}{3} \text{ of } 1\frac{1}{3} \div \frac{1}{3}$$

$$= \left[\frac{11}{3} - \frac{52}{9} \times 3 \times \frac{15}{13} \right]$$

$$= \frac{2}{3} \times \frac{4}{3} \times 3$$

$$= \frac{11 - 60}{3} \times \frac{3}{8} = -6\frac{1}{8}$$

Ex: Evaluate: $[7 + 7 \times (7 + 7 \div 7)] + 7 \div 7$

Sol: Using concept of BODMAS

$$= [7 + 7 \times (7 + 7 \div 7)] + 7 \div 7$$

$$= [7 + 7 \times (7 + 1)] + 1$$

$$= 63 + 1 = 64$$

Ex: The simplified value of the expression

$$1.0025 + 6.25 \times 10^{-6}$$

$$0.0025 + 0.95$$

Sol:

$$= \frac{1.0025 + 6.25 \times 10^{-6}}{0.0025 + 0.95}$$

$$= \frac{1.0025 + 0.00000625}{0.0025 + 0.95}$$

$$= \frac{1.00250625}{0.9525} = 1.0525$$

Equations based

Ex: What is 12% of 4% of 7% of 2×10^6 ?

$$\text{Sol: } 10^6 = (10^2)^3$$

$$\text{Number} = 2 \times 100^3 \times \frac{12}{100} \times \frac{4}{100} \times \frac{7}{100}$$

$$= 2 \times 100 \times 100 \times 100 \times \frac{12}{100} \times \frac{4}{100} \times \frac{7}{100}$$

$$= 2 \times 12 \times 4 \times 7 = 672$$

Ex: Simplify the following expression

$$25 - [16 - \{14 - (18 - \overline{8 + 3})\}]$$

Sol: Using concept of BODMAS

$$= 25 - [16 - \{14 - (18 - \overline{8 + 3})\}]$$

$$= 25 - [16 - \{14 - (18 - 11)\}]$$

$$= 25 - [16 - \{14 - 7\}] = 25 - [16 - 7]$$

$$= 25 - 9 = 16$$